

## OBSTETRICS

# The effect of maternal sleep-disordered breathing on the infant's neurodevelopment

Riva Tauman, MD; Luba Zuk, PhD; Shimrit Uliel-Sibony, MD; Jessica Ascher-Landsberg, MD; Shlomit Katsav, BSc; Mira Farber, BSc; Yakov Sivan, MD; Haim Bassan, MD

**OBJECTIVE:** We sought to examine the effect of maternal sleep-disordered breathing (SDB) on infant general movements (GMs) and neurodevelopment.

**STUDY DESIGN:** Pregnant women with uncomplicated full-term pregnancies and their offspring were prospectively recruited from a community and hospital low-risk obstetric surveillance. All participants completed a sleep questionnaire on second trimester and underwent ambulatory sleep evaluation (WatchPAT; Itamar Medical, Caesarea, Israel). They were categorized as SDB (apnea hypopnea index  $>5$ ) and controls. Infant GMs were assessed in the first 48 hours and at 8-11 and 14-16 weeks of age. At 12 months of age the Infant Developmental Inventory and the Brief Infant Sleep Questionnaire were administered.

**RESULTS:** In all, 74 women and their full-term infants were studied. Eighteen (24%) women had SDB. Mean birthweight was  $3347.1 \pm$

423.9 g. Median Apgar score at 5 minutes was 10 (range, 8–10). In adjusted comparisons, no differences were found between infants born to mothers with SDB and controls in GM scores in all 3 evaluations. Low social developmental score was detected at 12 months in 64% of infants born to SDB mothers compared to 25% of infants born to controls (adjusted  $P = .036$ ; odds ratio, 16.7). Infant snoring was reported by 41.7% of mothers with SDB compared to 7.5% of controls ( $P = .004$ ).

**CONCLUSION:** Our preliminary results suggest that maternal SDB during pregnancy has no adverse effect on neonatal and infant neuromotor development but may affect social development at 1 year.

**Key words:** fetal outcome, neurodevelopment, pregnancy, sleep-disordered breathing

Cite this article as: Tauman R, Zuk L, Uliel-Sibony S, et al. The effect of maternal sleep-disordered breathing on the infant's neurodevelopment. *Am J Obstet Gynecol* 2015;212:x.ex-x.ex.

Pregnancy is associated with significant changes in sleep with a large proportion of pregnant women experiencing some form of sleep disruption. Physiologic and hormonal changes that occur during pregnancy, particularly during the third trimester, place women at risk for developing sleep-disordered breathing (SDB).<sup>1,2</sup> Indeed, SDB is common during pregnancy and self-reported snoring has been observed in up to 46% of pregnant women.<sup>3-6</sup> The term “sleep-disordered breathing” refers

to a spectrum of abnormal respiration during sleep that ranges from primary (habitual) snoring to obstructive sleep apnea syndrome. It is characterized by episodic complete or partial obstruction of the airway during sleep, disruption of normal ventilation, intermittent hypoxemia, and sleep fragmentation.

It has been suggested that SDB during pregnancy may adversely influence the maternal and fetal well-being. Specifically, associations between SDB during pregnancy and gestational

diabetes, hypertension and pre-eclampsia,<sup>7-15</sup> fetal growth restriction, prematurity, cesarean delivery, and low Apgar scores have been reported.<sup>15-22</sup> However, these observations have thus far been inconclusive, one of the reasons being that most of the published literature is based on subjective assessment of sleep and lacks objective sleep measures.

Despite this inconsistency, in a recent study we found that maternal snoring was associated with enhanced fetal erythropoiesis and elevated cord interleukin 6.<sup>23</sup> We therefore hypothesized that maternal SDB with the resultant subtle intermittent hypoxia could influence the developing fetal brain.

To our knowledge, there are no reports on the effect of maternal SDB on infant neurodevelopmental status despite the ostensibly large body of data regarding the association of pediatric SDB with attention deficit, hyperactivity, and emotional and behavioral disturbances.<sup>24-31</sup> For all these reasons, we designed a prospective study using an

From the Sleep Disorders Center (Drs Tauman and Sivan, Ms Katsav, and Ms Farber), Child Neurology and Developmental Unit (Drs Uliel-Sibony and Bassan), and Departments of Physical Therapy (Dr Zuk) and Obstetrics and Gynecology (Dr Landsberg), Tel Aviv Medical Center; and Sackler School of Medicine, Tel Aviv University (Drs Tauman, Zuk, Uliel-Sibony, Landsberg, Sivan, and Bassan), Tel Aviv, Israel.

Received Sept. 26, 2014; revised Dec. 10, 2014; accepted Jan. 5, 2015.

This research was supported by grant number 707/12, Israel Science Foundation.

The authors report no conflict of interest.

Presented at the 26th annual meeting of the Associated Professional Sleep Societies, Baltimore, MD, June 1-5, 2013.

Corresponding author: Riva Tauman, MD. [tauman@tlvmc.gov.il](mailto:tauman@tlvmc.gov.il)

0002-9378/\$36.00 • © 2015 Elsevier Inc. All rights reserved. • <http://dx.doi.org/10.1016/j.ajog.2015.01.001>

objective sleep study measure to investigate the effects of maternal SDB on their infants. We hypothesized that infants born to mothers with SDB will exhibit lower developmental scores.

Our aims were to determine the effect of maternal SDB on neonatal and infant spontaneous general movements (GMs) and developmental outcome at 1 year.

## MATERIALS AND METHODS

Women in the third trimester of a singleton, uncomplicated pregnancy who attended low-risk obstetric surveillance from March 2009 through March

2012 were recruited. All participants completed a sleep questionnaire during the second trimester and underwent a sleep study during the third trimester of pregnancy. Infant GMs were assessed during the first 4 months of life. Development and sleep questionnaires were administered at 12 months. The study design and recruitment flowchart are presented in the [Figure](#).

The study was approved by the institutional review board. The study was registered at [ClinicalTrials.gov](#) (NCT00931099). Informed consent was obtained from all participants.

## Maternal and infant data

All women completed a sleep questionnaire (gestational week 25-27) regarding the presence of habitual snoring before and during pregnancy. Habitual snoring was defined as snoring at least 3 nights per week.<sup>32,33</sup> Pregnancy-onset snoring was considered present when habitual snoring began during pregnancy. All participants underwent an ambulatory overnight sleep study between 33-36 weeks of gestation using a validated ambulatory sleep technology (WatchPAT 200; Itamar Medical, Caesarea, Israel).<sup>34-37</sup> Apnea hypopnea index, respiratory disturbance index, oxygen desaturation index, mean oxygen saturation (SpO<sub>2</sub>) and SpO<sub>2</sub> nadirs were retrieved as previously described.<sup>37</sup> Women with an apnea hypopnea index >5 per hour of sleep were considered to have SDB.<sup>38</sup> We used the standard adult criteria for SDB, since there is no distinctive threshold for SDB in pregnancy. Medical records review was conducted by one of the researchers blinded to the sleep study results. Pertinent demographic (sex, gestational age, birthweight) and clinical (mode of delivery, Apgar scores at 1 and 5 minutes, and any perinatal complications) information were collected.

## Outcome data


### Spontaneous GMs assessment of infants

GMs were videotaped (15 minutes in length) for offline assessment in the first 48 hours of life, between 8-11 weeks of age, and between 14-16 weeks of age postterm. All infants were dressed in a diaper only, and were in the supine position. During off-time replay of the 3 periods videotaped, the GM quality was analyzed and assessed by an independent trained observer and the data scored. GMs were scored when infants were in active wakefulness. Periods of crying, fussing, and sucking were excluded from analysis.<sup>39,40</sup> The observer was unaware of the infant's group assignment (maternal SDB yes/no) as well as of other test results and clinical data.

### Sleep evaluation at the age of 1 year

Sleep was evaluated at 1 year of age using the Brief Infant Sleep Questionnaire

**FIGURE**  
**Recruitment flowchart**



Tauman. Maternal sleep-disordered breathing and child's neurodevelopment. *Am J Obstet Gynecol* 2015.

(BISQ). The BISQ is a validated sleep questionnaire assessing the infant's typical sleep patterns based on parental reports.<sup>41</sup> A yes/no question regarding snoring at the age of 1 year was added to the BISQ.

### Socioeconomic status

The modified, 2-factor index Hollingshead scale was used for determining the infant's socioeconomic status. The maternal and paternal scores were averaged.<sup>42</sup>

### Developmental questionnaire at the age of 1 year

At 1 year of age the developmental level was evaluated using the Infant Developmental Inventory questionnaire.<sup>43,44</sup> This

screening instrument is a parental report of the infant's developmental skills in 5 areas (gross motor, fine motor, social, self-help, and language). The instrument shows age-appropriate standards from which an estimated developmental age was recorded for each subdomain. An estimated developmental age, divided by the infant's chronological age, and multiplied by 100, yielded the subdomain developmental quotient. Averaging all 5 subdomains yields a total score for general developmental function, also expressed as developmental quotient.

### Statistical analysis

The statistical analyses were performed with software (SPSS, version 19.0; IBM

Corp, Armonk, NY). Comparisons of variables were conducted between infants born to women with SDB (study group) and infants born to non-SDB mothers (controls). Comparisons of variables according to group assignment were performed with independent *t* tests for continuous variables, and the  $\chi^2$  analyses for categorical variables. Linear and logistic regression analyses were performed for several dependent variables and were adjusted for the following confounders: gestational age, birthweight, sex, 5-minute Apgar score, and socioeconomic status. All reported *P* values are 2-tailed with statistical significance set at *P* < .05.

### RESULTS

In all, 74 healthy women completed a sleep questionnaire and underwent ambulatory sleep evaluation. Infant spontaneous GMs and 1-year developmental inventory were administered to 62 and 51 subjects, respectively (Figure). We were unable to obtain complete outcome data in all infants due to unavailability of parents or investigators. The demographic characteristics of the 23 children with a missing 1-year developmental inventory were similar to those with a complete developmental questionnaire (mean gestational age:  $39.4 \pm 1.3$  vs  $38.9 \pm 1.3$ , *P* = .14, and mean birthweight:  $3391.3 \pm 393.4$  vs  $3284.8 \pm 464.4$  g, *P* = .35). No participants had maternal or pregnancy-related complications such as hypertension, gestational diabetes, infection, or fetal growth restriction. Twenty-nine (39%) women reported habitual snoring. Ten (13%) were habitual snorers before and during pregnancy and 19 (26%) were pregnancy-onset snorers. Eighteen (24%) women met our criteria for SDB and 56 served as a control group. Of the 18 women with SDB, 5 did not report snoring, 4 reported snoring before and during pregnancy, and 9 reported pregnancy-onset snoring by questionnaire. Except for 2 premature deliveries at 36 weeks of gestation, all infants were born at term. In all, 45 (61%) newborns were male. Comparison of maternal and newborn characteristics as well as maternal sleep measures between the

TABLE 1

#### Comparison of maternal and newborns characteristics and maternal sleep measures of study group (sleep-disordered breathing) and controls

Characteristic	SDB (n = 18)	Controls (n = 56)	P value
<b>Maternal</b>			
Age, y	33.1 $\pm$ 3.7	32.7 $\pm$ 4.6	.78
BMI prepregnancy	25.3 $\pm$ 4.3	21.9 $\pm$ 2.5	.005
<b>Delivery</b>			
Normal	14 (78%)	42 (75%)	.95
Cesarean	3 (17%)	11 (20%)	
Instrumental	1 (6 %)	3 (5%)	
Gravida, median (range)	2.0 (1.0–7.0)	2.0 (1.0–5.0)	.84
Para, median (range)	2.0 (1.0–3.0)	1.0 (1.0–3.0)	.13
Total sleep time, min	334.4 $\pm$ 58.0	344.0 $\pm$ 83.0	.60
Respiratory disturbance index	14.4 $\pm$ 5.7	6.1 $\pm$ 3.2	< .0001
Apnea hypopnea index	11.2 $\pm$ 5.1	1.3 $\pm$ 1.4	< .0001
Oxygen desaturation index	3.1 $\pm$ 1.9	0.3 $\pm$ 0.4	< .0001
Mean SpO <sub>2</sub>	94.7 $\pm$ 0.9	96.1 $\pm$ 0.9	< .0001
SpO <sub>2</sub> nadir	89.6 $\pm$ 2.3	93.1 $\pm$ 2.2	< .0001
<b>Newborn</b>			
Male sex	10 (56%)	35 (63%)	.64
Apgar 1, median (range)	9 (4–9)	9 (3–9)	.14
Apgar 5, median (range)	10 (8–10)	10 (9–10)	.85
Gestational age, wk	39.3 $\pm$ 1.4	39.2 $\pm$ 1.4	.13
Birthweight, g	3371.0 $\pm$ 454.6	3356.7 $\pm$ 419.9	.9
Cord PH	7.28 $\pm$ 0.08	7.29 $\pm$ 0.09	.81

BMI, body mass index; SDB, sleep-disordered breathing; SpO<sub>2</sub>, oxygen saturation.

Tauman. Maternal sleep-disordered breathing and child's neurodevelopment. *Am J Obstet Gynecol* 2015.

TABLE 2

**Infant assessment of spontaneous general movements in first 48 hours, and at 8-11 and 14-16 weeks of age**

GM	Maternal SDB (n = 17)	Controls (n = 45)	P value	P value <sup>a</sup>
First 48 h	14.2 ± 3.9	15.0 ± 3.7	.47	.27
8-11 wk	22.5 ± 7.8	22.8 ± 7.3	.99	.98
14-16 wk	22.9 ± 8.1	23.8 ± 6.4	.66	.37
Total score	19.1 ± 7.3	20.5 ± 5.5	.56	.19

GM, general movement; SDB, sleep-disordered breathing.

<sup>a</sup> Derived from linear regression analysis adjusted for gestational age, birthweight, sex, and Apgar score at 5 min.Tauman. Maternal sleep-disordered breathing and child's neurodevelopment. *Am J Obstet Gynecol* 2015.

study group (SDB) and the controls are presented in Table 1.

**General movements**

The 3 sequential assessments of GMs were administered to 62 infants. No significant differences were found between the 2 groups before and after adjustment (Table 2).

**Outcome at 12 months**

The Infant Developmental Inventory questionnaire was administered at 1 year

of age to 51 subjects (11 from the SDB group and 40 from the control group). As shown in Table 3, no significant differences were found in the gross and fine motor, language, and self-help developmental scores between the 2 groups. Of note, the mean social developmental scores were lower in the study group when compared to the controls and almost reached statistical significance ( $97.8 \pm 19.7$  vs  $114.9 \pm 28.3$ ,  $P = .067$ ). A social developmental score  $<100$  was detected in 7 (64%) infants born to SDB

mothers compared to 10 (25%) controls (adjusted  $P = .036$ ; odds ratio, 16.7).

Sleep measures derived from the BISQ at the age of 1 year are presented in Table 4. No significant differences in nocturnal sleep duration, daytime sleep, number of nocturnal awakenings, sleep latency and wake after sleep onset, and frequency of problematic sleep were found between children born to mothers with SDB compared with controls. However, infant snoring was reported by 5 of 12 (41.7%) mothers with SDB compared to 3 of 40 (7.5%) controls ( $P = .004$ ).

**COMMENT**

This is the first study that investigated the effect of maternal SDB during pregnancy on neonatal and infant neurodevelopment. Our preliminary findings suggest that maternal SDB has no impact on neonatal and infant spontaneous GM. Nevertheless, despite the similar early neuromotor status, assessment at 1 year suggested that infants born to mothers with SDB exhibited lower social

TABLE 3

**Developmental scores at age of 1 year**

Variable	SDB (n = 11)	Controls (n = 40)	P value	P value <sup>a</sup>	OR
Age, mo	12.9 ± 1.6	13.0 ± 1.4	.88		
Male sex	5 (45%)	24 (60%)	.34		
Parental socioeconomic status	24.2 ± 2.8	26.9 ± 7.5	.36		
Social score	97.8 ± 19.7	114.9 ± 28.3	.067	.13	
Social score $<100$ , n (%)	7 (64)	10 (25)	.029	.036	16.7
Self-help score	115.3 ± 9.7	115.2 ± 21.0	.93	.63	
Self-help score $<100$ , n (%)	2 (18)	0 (0)	.13	.99	
Gross motor score	114.7 ± 31.7	115.2 ± 21.0	.95	.43	
Gross motor score $<100$ , n (%)	4 (36)	8 (20)	.28	.11	4.5
Fine motor score	115.6 ± 21.2	115.2 ± 21.1	.95	.59	
Fine motor score $<100$ , n (%)	1 (9)	7 (18)	.46	.54	0.47
Language score	114.7 ± 21.6	117.0 ± 19.2	.74	.57	
Language score $<100$ , n (%)	3 (27)	5 (13)	.27	.15	6.2
Total developmental score	111.6 ± 17.4	114.3 ± 18.2	.66	.34	
Total score $<100$ , n (%)	3 (27)	6 (15)	.39	.13	6.3

OR, odds ratio; SDB, sleep-disordered breathing.

<sup>a</sup> Derived from linear regression analysis adjusted for gestational age, birthweight, sex, Apgar score at 5 min, and parental socioeconomic status.Tauman. Maternal sleep-disordered breathing and child's neurodevelopment. *Am J Obstet Gynecol* 2015.



TABLE 4

**Brief Infant Sleep Questionnaire results at age of 1 year in infants born to mothers with sleep-disordered breathing and to controls**

Variable	SDB (n = 12)	Controls (n = 40)	P value
Nocturnal sleep time, min	630.0 ± 54.3	594.4 ± 59.8	.07
Daytime sleep, min	146.3 ± 33.9	139.9 ± 43.6	.64
No. of night awakenings	1.83 ± 1.1	1.7 ± 1.4	.69
Wakefulness time after sleep onset, min	17.9 ± 16.6	22.1 ± 35.7	.7
Sleep latency, min	16.3 ± 7.7	21.7 ± 16.4	.14
Bedtime	8.06 pm ± 0.27	8.08 pm ± 0.36	.85
You consider your child's sleep to be a problem, n (%)	3 (25)	9 (22.5)	.85

SDB, sleep-disordered breathing.

Tauman. Maternal sleep-disordered breathing and child's neurodevelopment. *Am J Obstet Gynecol* 2015.

developmental scores and increased rate of snoring compared to controls.

To test our hypothesis on the effects of maternal SDB on the developing brain we performed 3 sequential evaluations of the neonatal and infant GMs and found similar scores in the 2 groups. We chose the GM assessment because of its good predictive ability of neurodevelopmental outcome.<sup>45</sup> Spontaneous GM evaluation is primarily based on neuromotor functions; therefore future studies should use behaviorally oriented evaluations to further test this hypothesis.

At 12 months of age, we administered a developmental questionnaire and found lower social developmental scores in infants born to pregnant mothers with SDB while motor and speech development were not different between the 2 groups.

Our finding supports previous observation of Hermans et al,<sup>46</sup> who demonstrated a significantly decreased grooming behavior and alterations in passive avoidance and locomotor activity in rat pups born of dams exposed to intermittent antenatal hypoxia. In their study, the abnormal neonatal behavior disappeared at postnatal day 30. Most neonatal neuromotor reflexes and Morris water maze performance (learning and memory testing) at 29 days were not affected by intermittent pregnancy hypoxia.<sup>46</sup> Gozal et al,<sup>47</sup> in their study of gestational intermittent hypoxia in a rat model, also found no effects on Morris water maze testing at postnatal day 30.

In a neurohistological study of intermittent gestational hypoxia in a rat model, Zechel et al<sup>48</sup> reported a transient delay in neuronal migration associated with alterations of multiple proteins early in the postpartum period. Based on these animal studies one could hypothesize that intermittent gestational hypoxia, increased intrathoracic pressure swings, or sleep fragmentation associated with maternal SDB could alter fetal brain developmental processes, either directly or via mediators. For example, enhanced fetal hypothalamic–pituitary–adrenal activity was found in a rat model of intermittent gestational hypoxia resulting in anxiety-like behavior in adult male offspring.<sup>49</sup> Other possible mediators may involve maternal and fetal perturbation of the autonomic nervous system<sup>50</sup> and exposure of the fetus to inflammatory mediators,<sup>23</sup> which ultimately may affect fetal brain development.

Nevertheless, comparison between human and animal models of maternal SDB should be viewed cautiously since the level of hypoxia applied in animal models may exceed that of human models.

Our finding of increased rate of snoring (41.7%) among infants born to mothers with SDB is in agreement with previous publications showing that family history of SDB is a risk factor for developing SDB during childhood.<sup>51</sup> The contribution of other risk factors such as allergy, adenotonsillar hypertrophy, and weight was not investigated in the present

study. However, this study is the first to show that maternal SDB during pregnancy increases the risk for SDB in the infant and, more importantly, that snoring is already present at 1 year of age.

Given the known association between pediatric SDB and behavioral emotional impairments, attention difficulties, impulsivity, hyperactivity, and daytime sleepiness, it is possible that our findings of delayed social development in infants born to mothers with SDB are also mediated by the infants' SDB.<sup>24-31</sup>

Based on the nature of the neuro-behavioral consequence of SDB,<sup>52</sup> early screening and intervention in children born to snoring mothers may potentially attenuate the progression of the neuro-cognitive outcomes.

This study has several limitations: our cohort consists of a relatively small number of mothers with SDB and includes some infants with incomplete outcome data that limit the ability to conduct subgroup analysis between chronic and pregnancy-onset SDB. In addition, the infants' development and sleep habits were based on a parental questionnaire and not on objective testing. Therefore, our results should be viewed as preliminary. Future studies should implement comprehensive neonatal behavioral assessments and a formal developmental outcome instrument that focuses on infants' communication. Notwithstanding such considerations, the strength of our study is its prospective longitudinal

follow-up design, utilizing a formal objective sleep study measure in a homogenous low-risk pregnancy cohort.

In conclusion, this is the first study examining the effect of gestational maternal SDB on the neonatal and infant neurodevelopment. Our preliminary results suggest that maternal SDB during pregnancy has no large adverse effect on neonatal and infant neuromotor development but it may affect social development at the age of 1 year. In addition, maternal SDB is associated with increased rate of infant snoring. Further studies using comprehensive developmental tools and larger sample size are warranted. ■

#### ACKNOWLEDGMENTS

We wish to thank the pregnant women and their children who participated in this study.

#### REFERENCES

- Pien GW, Schwab RJ. Sleep disorders during pregnancy. *Sleep* 2004;27:1405-17.
- Izci B, Riha RL, Martin SE, et al. The upper airway in pregnancy and pre-eclampsia. *Am J Respir Crit Care Med* 2003;167:137-40.
- Leung PL, Hui DSC, Leung TN, Yuen PM, Lau TK. Sleep disturbances in Chinese pregnant women. *BJOG* 2005;112:1568-71.
- Loube DI. Self-reported snoring in pregnancy: association with fetal outcome. *Chest* 1996;109:885-9.
- Franklin KA. Snoring, pregnancy-induced hypertension and growth retardation of the fetus. *Chest* 2000;117:137-41.
- Ursavas A, Karadag M, Nalci N, Ercan I, Gozu RO. Self-reported snoring, maternal obesity and neck circumference as risk factors for pregnancy-induced hypertension and pre-eclampsia. *Respiration* 2008;76:33-9.
- Qiu C, Enquobahrie D, Frederick I, Abetew D, Williams M. Glucose intolerance and gestational diabetes risk in relation to sleep duration and snoring during pregnancy: a pilot study. *BMC Womens Health* 2010;10:17.
- Bourjeily G, Raker CA, Chalhoub M, Miller MA. Pregnancy and fetal outcomes of symptoms of sleep-disordered breathing. *Eur Respir J* 2010;36:849-55.
- Facco FL, Grobman WA, Kramer J, Ho KH, Zee PC. Self-reported short sleep duration and frequent snoring in pregnancy: impact on glucose metabolism. *Am J Obstet Gynecol* 2010;203:142.e1-5.
- Facco FL, Lappen J, Lim C, Zee PC, Grobman WA. Preeclampsia and sleep-disordered breathing: a case-control study. *Pregnancy Hypertens* 2013;3:133-9.
- Louis J, Auckley D, Miladinovic B, et al. Perinatal outcomes associated with obstructive sleep apnea in obese pregnant women. *Obstet Gynecol* 2012;120:1085-92.
- Blyton DM, Skilton MR, Edwards N, Hennessy A, Celemajer DS, Sullivan CE. Treatment of sleep disordered breathing reverses low fetal activity levels in preeclampsia. *Sleep* 2013;36:15-21.
- O'Brien LM, Bullough AS, Owusu JT, et al. Pregnancy-onset habitual snoring, gestational hypertension, and preeclampsia: prospective cohort study. *Am J Obstet Gynecol* 2012;207:487.e1-9.
- Yinon D, Lowenstein L, Suraya S, et al. Preeclampsia is associated with sleep-disordered breathing and endothelial dysfunction. *Eur Respir J* 2006;27:328-33.
- Louis JM, Mogos MF, Salemi JL, Redline S, Salihu HM. Obstructive sleep apnea and severe maternal-infant morbidity/mortality in the United States, 1998-2009. *Sleep* 2014;37:843-9.
- Yin TT, Williams N, Burton C, et al. Hypertension, fetal growth restriction and obstructive sleep apnea in pregnancy. *Eur J Obstet Gynecol Reprod Biol* 2008;141:35-8.
- Ayrim A, Keskin EA, Ozol D, Onaran Y, Yildirim Z, Kafali H. Influence of self-reported snoring and witnessed sleep apnea on gestational hypertension and fetal outcome in pregnancy. *Arch Gynecol Obstet* 2011;283:195-9.
- Tauman R, Sivan Y, Katsav S, Greenfeld M, Many A. Maternal snoring is not associated with fetal growth restriction. *J Matern Fetal Neonatal Med* 2011;25:1283-6.
- Franklin KA, Holmgren PA, Jonsson F, Poromaa N, Stenlund H, Svanborg E. Snoring, pregnancy-induced hypertension, and growth retardation of the fetus. *Chest* 2000;117:137-41.
- Bourjeily G, Chalhoub M, Miller MA. Pregnancy and fetal outcomes of symptoms of sleep disordered breathing. *Eur Respir J* 2010;36:849-55.
- Louis JM, Auckley D, Sokol RJ, Mercer BM. Maternal and neonatal morbidities associated with obstructive sleep apnea complicating pregnancy. *Am J Obstet Gynecol* 2010;202:261.e1-5.
- Chen YH, Kang JH, Lin CC, Wang IT, Keller JJ, Lin HC. Obstructive sleep apnea and the risk of adverse pregnancy outcomes. *Am J Obstet Gynecol* 2012;206:136.e1-5.
- Tauman R, Many A, Deutsch V, et al. Maternal snoring during pregnancy is associated with enhanced fetal erythropoiesis—a preliminary study. *Sleep Med* 2011;12:518-22.
- Ali NJ, Pitson D, Stradling JR. Natural history of snoring and related behavior problems between the ages 4 and 7 years. *Arch Dis Child* 1994;71:74-6.
- Biggs SN, Vlahandonis A, Anderson V, et al. Long-term changes in neurocognition and behavior following treatment of sleep disordered breathing in school-aged children. *Sleep* 2014;37:77-84.
- Landau YE, Bar-Yishay O, Greenberg-Dotan S, Goldbart AD, Tarasiuk A, Tal A. Impaired behavioral and neurocognitive function in preschool children with obstructive sleep apnea. *Pediatr Pulmonol* 2012;47:180-8.
- Kohler MJ, Lushington K, Kennedy JD. Neurocognitive performance and behavior before and after treatment for sleep-disordered breathing in children. *Nat Sci Sleep* 2010;16:159-85.
- Owens JA. Neurocognitive and behavioral impact of sleep disordered breathing in children. *Pediatr Pulmonol* 2009;44:417-22.
- Friedman BC, Hendeles-Amitai A, Kozminsky E, et al. Adenotonsillectomy improves neurocognitive function in children with obstructive sleep apnea syndrome. *Sleep* 2003;26:999-1005.
- Scullin MH, Ornelas C, Montgomery-Downs HE. Risk for sleep-disordered breathing and home and classroom behavior in Hispanic preschoolers. *Behav Sleep Med* 2011;9:194-207.
- Montgomery-Downs HE, Gozal D. Sleep-associated respiratory disorders and their psychobehavioral consequences in children. *Handb Clin Neurol* 2011;98:489-99.
- Kump K, Whalen C, Tishler PV, et al. Assessment of the validity and utility of a sleep-symptom questionnaire. *Am J Respir Crit Care Med* 1994;150:735-41.
- Young T, Shahar E, Nieto FJ, et al. Predictors of sleep-disordered breathing in community-dwelling adults: the Sleep Heart Health Study. *Arch Intern Med* 2002;162:893-900.
- Hedner J, White DP, Malhotra A, et al. Sleep stages based on autonomic signals: a multi-center validation study. *J Clin Sleep Med* 2011;7:301-6.
- Choi JH, Kim EJ, Kim YS, et al. Validation study of portable device for the diagnosis of obstructive sleep apnea according to the new AASM scoring criteria: Watch PAT 100. *Acta Otolaryngol* 2010;30:838-43.
- Herscovici S, Pe'er A, Pappan S, Lavie P. Detecting REM sleep from the finger: an automatic REM sleep algorithm based on peripheral arterial tonometry (PAT) and actigraphy. *Physiol Meas* 2007;28:129-40.
- Pittman SD, Ayas NT, MacDonald MM, Malhotra A, Fogel RB, White DP. Using a wrist worn device based on peripheral arterial tonometry to diagnose obstructive sleep apnea: in laboratory and ambulatory validation. *Sleep* 2004;27:923-33.
- American Academy of Sleep Medicine Task Force. Sleep-related breathing disorders in adults: recommendation for syndrome definition and measurement techniques in clinical research. *Sleep* 1999;22:667-89.
- Hadders-Algra M. General movements: a window for early identification of children at high risk for developmental disorders. *J Pediatr* 2004;145(Suppl):S12-8.

40. Einspieler C, Prechtl HF. Prechtl's assessment of general movements: a diagnostic tool for the functional assessment of the young nervous system. *Ment Retard Dev Disabil Res Rev* 2005;11:61-7.
41. Sadeh A. A brief screening questionnaire for infant sleep problems: validation and findings for an Internet sample. *Pediatrics* 2004;113:e570-7.
42. Freedman LZ, Hollingshead AB. Neurosis and social class, I: social interaction. *Am J Psychiatry* 1957;113:769.e75.
43. Creighton DE, Sauve RS. The Minnesota Infant Developmental Inventory in the developmental screening of high-risk infants at 8 mo. *Can J Behav Sci* 1988;20:424-33.
44. Eaves RC, Reilly A. Factor analysis of the Minnesota Infant Developmental Inventory based on a Hispanic migrant population. *Educ Psychol Meas* 2000;60:271-85.
45. Zuk L. Fetal and infant spontaneous general movements as predictors of developmental disabilities. *Dev Disabil Res Rev* 2011;17: 93-101.
46. Hermans RH, Hunter DE, McGivern RF, Cain CD, Longo LD. Behavioral sequelae in young rats of acute intermittent antenatal hypoxia. *Neurotoxicol Teratol* 1992;14: 119-29.
47. Gozal D, Reeves SR, Row BW, Neville JJ, Guo SZ, Lipton AJ. Respiratory effects of gestational intermittent hypoxia in the developing rat. *Am J Respir Crit Care Med* 2003;167: 1540-7.
48. Zechel JL, Gamboa JL, Peterson AG, Puchowicz MA, Selman WR, Lust WD. Neuronal migration is transiently delayed by prenatal exposure to intermittent hypoxia. *Birth Defects Res Dev Reprod Toxicol* 2005;74:287-99.
49. Fan JM, Chen XQ, Jin H, Du JZ. Gestational hypoxia alone or combined with restraint sensitizes the hypothalamic-pituitary-adrenal axis and induces anxiety-like behavior in adult male rat offspring. *Neuroscience* 2009;159:1363-73.
50. Hakim F, Gozal D, Kheirandish-Gozal L. Sympathetic and catecholaminergic alterations in sleep apnea with particular emphasis on children. *Front Neurol* 2012;3:1-13.
51. Alexopoulos EI, Charitos G, Malakasioti G, et al. Parental history of adenotonsillectomy is associated with obstructive sleep apnea severity in children with snoring. *J Pediatr* 2014;164: 1352-7.
52. Jackman AR, Biggs SN, Walter LM, et al. Sleep-disordered breathing in preschool children is associated with behavioral, but not cognitive, impairments. *Sleep Med* 2012;13:621-31.